

# *Suicide Among Veterans and Other Americans*

## **2001–2014**

Office of Suicide Prevention

**3 August 2016**



U.S. Department  
of Veterans Affairs

# Contents

I. Introduction.....	3
II. Executive Summary.....	4
III. Background .....	5
IV. Methodology .....	5
V. Results – Part 1: Suicide Among VHA Patients With Comparison to the U.S. General Population, 2001–2014.....	6
A. VHA Patient Population.....	6
B. Mental Health and Substance Use Disorders Among VHA Patients.....	8
C. History of Non-Fatal Suicide Attempt .....	11
D. Total and Gender-Specific VHA Suicide Rates.....	14
E. Differences in VHA Patient Suicide Rates by Age and Gender .....	16
F. Comparison of Suicide Rates among VHA Users and the General U.S. Population .....	17
G. Suicide Among OEF/OIF/OND VHA Users .....	19
VI. Results – Part 2: Suicide Among All U.S. Veterans, 2001–2014.....	20
A. Magnitude of Veteran Suicide Mortality .....	21
B. Comparison of Suicide Rates Among Veterans Who Do and Do Not Use VHA Services, 2001–2014.....	23
C. Comparison of Veteran and Adult Civilian Suicide Risk, 2001–2014 .....	24
D. Mechanisms of Veteran and Civilian Suicide, 2001–2014 .....	38
E. Understanding the Burden of Veteran Suicide: Magnitude vs. Risk.....	41
VII. Summary and Discussion of Findings.....	45
VIII. Suicide Data Report Part II.....	46

## I. Introduction

The top priority of the U.S. Department of Veterans Affairs (VA) is the health and well-being of all of our nation's Veterans, including Veterans who have chosen not to enroll in, or are not eligible for, VA health care. As the largest integrated health care system in the country, VA is committed to providing timely access to high-quality, recovery-oriented mental health care that anticipates and responds to Veterans' needs, such as treatment for PTSD, substance use disorders, depression, and suicidal ideation.

In 2014, suicide was the 10<sup>th</sup> leading cause of death in the United States and rates of suicide in the U.S. general population are increasing. Centers for Disease Control and Prevention (CDC) data released in an [April 2016 report](#) indicated that between 1999 and 2014, suicide rates increased among the general population, for both males and females and for all ages. Regardless of suicide rates or the number of cases, one life lost to suicide is too many.

VA has worked tirelessly to develop suicide prevention resources for every Veteran who is experiencing a mental health crisis, whether or not that Veteran is enrolled in the VA Health Care System. In fact, of about 21.6 million\* Veterans across the country — including almost 2 million women — just over 8.5 million\*\* are enrolled for care from a VA provider.

VA is committed to identifying and reaching all Veterans who may be at risk for suicide and continues to enhance programs designed to reduce risk among those who receive services from the Veterans Health Administration (VHA). As highlighted in a recent VA-led Call to Action to Prevent Veteran Suicide, eliminating the burden of suicide among Veterans will require participation from a broad group of federal government and community partners. In recognition of this need, VA and its partners are developing innovative strategies to find and help Veterans at risk for suicide through community-based collaborations and expanded supportive services.

As part of the Call to Action, VA has undertaken the most comprehensive analysis of Veteran suicide in our nation's history, examining more than 55 million Veteran records from 1979 to 2014 from all 50 states and four territories. This report describes the results of this effort. It builds on data from previous VA Suicide Data Reports, which were primarily limited to information on Veterans who used VHA health services or from mortality records obtained directly from a small number 20 of states and approximately 3 million records.

This report on Veteran suicide is unprecedented in its breadth and depth of information about the characteristics of suicide among Veterans. It contains the first comprehensive assessment of differences in rates of suicide among Veterans with and without use of VHA services and comparisons between Veterans and other Americans. This report serves as a foundation for informing and evaluating suicide prevention efforts inside the VHA health care system and for developing lifesaving collaborations with community health care partners.

\*Source: VA Benefits & Health Care Utilization Pocket Card, Updated 5/13/16; Veteran Population as of 09/30/15 (<http://www.va.gov/vetdata/docs/pocketcards/fy2016q3.pdf>)

\*\*Source: VA Benefits & Health Care Utilization Pocket Card, Updated 5/13/16; Produced by the National Center for Veterans Analysis and Statistics. (<http://www.va.gov/vetdata/docs/pocketcards/fy2016q3.pdf>)

## II. Executive Summary

This report provides information regarding suicide mortality for the years 2001–2014. It incorporates the most recent mortality data from the VA/Department of Defense (DoD) Joint Suicide Data Repository and includes information for deaths from suicide among all known Veterans of U.S. military service. Data for the Joint VA/DoD Suicide Data Repository were obtained from the National Center for Health Statistics' National Death Index through collaboration with the DoD, the CDC, and the VA/DoD Joint Suicide Data Repository initiative. Data available from the National Death Index include reports of mortality submitted from vital statistics systems in all 50 U.S. states, New York City, Washington D.C., Puerto Rico, and the U.S. Virgin Islands.

This report is unprecedented in its comprehensive analysis of suicide rates among all U.S. Veterans. Unlike previous Department reports, this report provides information on all recorded suicides among all Veterans living in the U.S. Additional enhancements include direct comparisons of Veterans' suicide rates with those of analogous civilian populations, calculation of suicide rates among populations with known elevations in suicide risk (i.e., mental health diagnoses), groups with emerging risk (i.e., patients prescribed opioids), and comparisons between Veterans with and without use of VHA services. In contrast to previous VA reports, rates of suicide have been calculated by calendar year to facilitate comparison with national statistics and reports from other agencies.

Findings on suicide counts and rates are based on analyses conducted at the VHA Office of Suicide Prevention with support from the VISN 2 Center of Excellence for Suicide Prevention, VISN 19 Mental Illness Research, Education and Clinical Care Center, and Post-Deployment Health Service. Results from analyses included in this report were obtained using all available information to identify Veterans who died by suicide. This report includes the years 2001–2014. Subsequent analyses will include data from earlier years.

Key findings from this year's report include:

- In 2014, an average of 20 Veterans died by suicide each day. Six of the 20 were users of VHA services.
- In 2014, Veterans accounted for 18 percent of all deaths by suicide among U.S. adults and constituted 8.5 percent of the U.S. adult population (ages 18+). In 2010, Veterans accounted for 20.2 percent of all deaths by suicide and represented 9.7 percent of the U.S. adult population.
- The burden of suicide resulting from firearm injuries remains high. In 2014, about 67 percent of all Veteran deaths by suicide were the result of firearm injuries.
- There is continued evidence of a high burden of suicide among middle-aged and older Veterans. In 2014, about 65 percent of all Veterans who died by suicide were age 50 or older.
- After adjusting for differences in age and gender, risk for suicide was 21 percent higher among Veterans when compared with U.S. civilian adults. (2014)
- After adjusting for differences in age, risk for suicide was 18 percent higher among male Veterans when compared with U.S. civilian adult males. (2014)
- After adjusting for differences in age, risk for suicide was 2.4 times higher among female Veterans when compared with U.S. civilian adult females. (2014)
- In 2014, rates of suicide were highest among younger Veterans (ages 18–29) and lowest among older Veterans (ages 60+). Furthermore, rates of suicide among Veterans age 70 and older were lower than rates of suicide among civilians in the same age group.

### III. Background

Rates of suicide have been increasing for both males and females and across all age groups in the United States. According to a recent CDC report, the age-adjusted rate of suicide increased by 24 percent between 1999 and 2014.<sup>1</sup> Findings from this same report show that increases in rates of suicide were higher between 2006 and 2014 than they were during earlier time periods. Despite having lower rates of suicide when compared to men in the general population, rates of suicide increased more among women than among men during the study period. While overall rates of suicide have increased in the United States, suicides resulting from a firearm injury have decreased since 1999. According to the CDC, the proportion of suicides resulting from a firearm injury decreased by more than 10 percent among men and 16 percent among women in the U.S. general population. Finally, different patterns were seen in the distribution of suicide rates across age groups for men and women. With some slight variability, rates of suicide increased with age among males in the U.S. general population, with the highest rates of suicide among males aged 75 years and older. In contrast, rates of suicide among women in the U.S. general population peaked during middle age, with the highest rates among women ages 45–64.

The VA has released two previous Suicide Data Reports (2012, 2014). While these previous reports did not include information on the characteristics of suicide among all Veterans, the available information did provide valuable insight into potential differences between suicide among those with history of U.S. military service and other Americans. Of particular importance were findings of increases in rates of suicide among younger Veterans (18–29 years of age), gender-based differences in changes in rates among women Veterans who used VHA services, and a comparatively high prevalence (approximately 66 percent) of suicides resulting from a firearm injury. Results included in this report provide the first systematic assessment of characteristics of suicide among Veterans with and without use of VHA services and comparison to rates of suicide among other Americans (i.e., civilians).

### IV. Methodology

Data for this report were obtained by linking information from VA and DoD administrative records with cause of death information included in CDC's National Death Index. Information from multiple program offices and record systems was combined to create a comprehensive population record of Veterans for the years of interest. From VA, information was obtained from population rosters maintained by the Office of Policy and Planning, from deployment and service rosters maintained by the Post-Deployment Health Service, and from VHA clinical and administrative records. Information on all Veterans who separated from active duty service or who had been activated during service in a Reserve component or the National Guard was obtained from the DoD Defense Manpower Data Center. In total, more than 50 million records were submitted to the NDI for retrieval of information on fact and cause of death.

This report is divided into two sections, parts A and B. The first section, Part A, includes information on rates of suicide among all VHA patients and compares these rates to suicide rates among the general U.S. population. Consistent with past practice, rates of suicide and estimates of relative risk presented in the first section include information on all users of VHA services (Veteran and other users) with comparison to rates of suicide among members of the U.S. general population. The second section, Part B, includes information on rates of suicide among only Veterans, including both those who used VHA clinical services and those who did not. The report compares these with suicide rates among adult civilians (i.e., excluding Veterans). In both sections of this report, Veteran suicide decedents were considered users of VHA services if there was at least one record of inpatient or outpatient care in the calendar year of or before death.

## **V. Results – Part 1: Suicide Among VHA Patients With Comparison to the U.S. General Population, 2001–2014**

This section provides information regarding suicide mortality among all patients, including those who were not Veterans, versus those who received VHA health care services. Findings on suicide counts, rates, and risk factors in this section are based on analyses conducted at the VHA's Office of Mental Health Operations' Serious Mental Illness Treatment Resource and Evaluation Center. For the years 2001–2014, the rate of suicide among patients who used VHA health care services in the year of death or in the previous calendar year have been evaluated overall and by gender and age group.


### **A. VHA Patient Population**

The VHA provides health care to a large and diverse patient population and, as is true with many health systems, provides care for patients with complex health problems, some of which are associated with increased risk for suicide. It is also important to note, that not all Veterans are equally eligible to receive VHA services. One way of understanding the characteristics of VHA's patient population is to examine the type of eligibility assigned to each Veteran who received VHA care. Veterans are assigned priority groups that determine their eligibility status for VHA services. Eligibility is largely, but not solely, based on service connected disability level and income. Since 2001, some notable changes in the VHA patient population have occurred. Specifically, the proportion of VHA Veterans with a 50 percent or higher service connected disability has more than doubled, increasing from 11.7 percent in 2001 to 25.9 percent of all VHA patients in 2014. At the same time, the proportion of non-Veteran patients has decreased significantly, dropping from 8.8 percent in 2001 to 4.4 percent of all VHA patients in 2014.<sup>2</sup> Additional changes in the patient population distribution by priority enrollment group can be seen in Figure 1.

1. Curtin SC, Warner M, Geedegaard H. Increase in suicide in the United States, 1999–2014. NCHS data brief, no. 241. Hyattsville, MD: National Center for Health Statistics. 2016.

2. Details on priority group enrollment criteria can be found at: [http://www.va.gov/HEALTHBENEFITS/resources/priority\\_groups.asp](http://www.va.gov/HEALTHBENEFITS/resources/priority_groups.asp)

Figure 1. VHA Patient Distribution by Enrollment Priority Group (percent), 2001 and 2014




Priority Group Descriptions: **Group 1** = ≥ 50 percent service connected disability; **Group 2** = 30 percent – 40 percent service connected disability; **Group 3** = 20 percent service connected disability, prisoners of war, other special categories; **Group 4** = Veterans who are receiving aid and attendance or housebound benefits from VA or Veterans who have been determined to be catastrophically disabled; **Group 5** = nonservice-connected Veterans and noncompensable service-connected Veterans rated 0 percent disabled by VA with annual income below the VA and geographically (based on your resident zip code) adjusted income limits, Veterans receiving VA pension benefits, Veterans eligible for Medicaid programs; **Group 6** = all other Veterans not required to make a co-pay; **Group 7** = Veterans with gross household income below the geographically adjusted income limits (GMT) for their resident location and who agree to pay copays; **Group 8** = Veterans with gross household income above the VA and the geographically adjusted income limits for their resident location and who agrees to pay copays.

## B. Mental Health and Substance Use Disorders Among VHA Patients

Mental health disorders, including major depression and other mood disorders, have been associated with increased risk for suicide.<sup>3</sup> Since 2001, the proportion of VHA users with mental health conditions or substance use disorders (SUD) has increased from approximately 27 percent in 2001 to more than 40 percent in 2014. The increased prevalence of mental health disorders among VHA patients when compared to the U.S. adult population should not be taken as an indicator of the overall mental health of the larger Veteran population. Rather, this information may explain differences in rates among VHA patients when compared to rates of suicide in the general population.

Figure 2. Percentage of VHA Users With Diagnoses of Mental Health (MH) Conditions/Substance Use Disorders (SUD) by Calendar Year




**Main Finding: The percentage of VHA users diagnosed with a mental health or substance use disorder has increased substantially since 2001.**

3. Harris, E.C. & Barraclough, B. (1997) Suicide as an outcome for mental health disorders. A meta-analysis. Br J Psychiatry, 170, 205-228.



Despite a growing patient population with known risk factors for suicide, rates of suicide remained stable among VHA patients diagnosed with a mental health condition or substance use disorder between 2001 and 2014 (Figure 3). Rates were highest among those diagnosed with severe depression (BPD), which as shown in Figure 2 represents less than 5 percent of VHA patients. In contrast, rates among those with any mental health condition or substance use disorder decreased from 77.6 to 57.0 per 100,000 between 2001 and 2014; for those diagnosed with depression, the rate decreased from 100.01 to 70.9 per 100,000 between 2001 and 2014.


Figure 3. Suicide Rate Per 100,000 Person-Years, Among VHA Users by Mental Health (MH) Conditions/Substance Use Disorders (SUD), by Condition and Calendar Year



**Main Finding: As compared to 2001, rates of suicide have decreased among VHA patients diagnosed with a mental health or substance use disorder.**

Figure 4 provides information on rates of suicide among those patients diagnosed with opioid use disorder (OUD), a condition with emerging evidence of suicide risk. As shown in Figure 4, rates of suicide have increased among VHA patients with OUD and are comparable to rates of suicide among VHA patients diagnosed with severe depression (BPD).

Figure 4. Suicide Rate Per 100,000 Person-Years Among VHA Users by Receipt of Opioid Use Disorder Diagnosis by Calendar Year



**Main Finding: Rates of suicide were elevated among VHA patients diagnosed with an opioid use disorder and have increased since 2001.**

Overall, suicide rates are highest among patients with mental health and substance use disorder diagnoses who are in treatment and lower among those who received a MH diagnoses but were not sick enough to require enhanced care from a MH provider (Table 1).

Table 1. Suicide Rates by Receipt of Mental Health (MH)/Substance Use Disorder (SUD) Diagnosis or Treatment and Calendar Year


Characteristics	Calendar Year													
	CY 01	CY 02	CY 03	CY 04	CY 05	CY 06	CY 07	CY 08	CY 09	CY 10	CY 11	CY 12	CY 13	CY 14
Total	39.9	38.9	34.9	35.9	34.9	35.9	35.1	38.4	37.0	36.3	38.9	37.9	38.8	39.0
With MH treatment	84.0	81.4	68.9	69.8	65.0	68.1	64.3	70.7	65.1	62.3	66.3	63.3	62.3	65.0
Without MH treatment	27.7	27.6	25.8	26.6	26.3	26.3	25.9	27.2	26.5	26.0	27.4	26.8	28.1	26.7
With MH/SUD diagnosis	77.6	74.1	63.5	62.1	59.4	59.0	57.6	60.4	57.6	55.2	58.2	56.1	55.8	56.5
Without MH/SUD diagnosis	24.7	24.5	22.3	23.3	22.5	23.6	22.5	25.4	24.2	24.1	25.9	25.3	26.7	26.3
<b>With MH/SUD diagnosis and MH treatment</b>	89.8	88.2	74.6	75.2	70.6	72.4	69.1	75.3	68.4	65.5	69.3	65.5	65.5	67.5
<b>With MH/SUD diagnosis, without MH treatment</b>	52.9	49.0	46.0	43.1	43.2	39.4	40.4	36.2	38.9	36.4	37.3	38.2	36.9	34.3
Without MH/SUD diagnosis, with MH treatment	38.7	27.9	23.1	24.6	18.3	30.7	21.7	32.0	36.9	35.4	42.8	46.9	39.8	47.2
<b>Without MH/SUD diagnosis, without MH treatment</b>	24.2	24.3	22.3	23.3	22.7	23.3	22.5	25.1	23.6	23.5	24.9	23.9	25.8	24.7

**Main Finding: VHA patients with diagnosis who accessed mental health treatment services have higher rates of suicide than other VHA patients.**

### C. History of Non-Fatal Suicide Attempt


A history of non-fatal suicide attempts is recognized to be among the most robust risk factors for suicide. Among VHA patients, reports of suicide attempt can be identified through review of external injury codes associated with health services (i.e., obtained from medical records) or from the Suicide Prevention Applications Network (SPAN), VHA’s internal suicide event case management and tracking system. As shown in Figure 5, monthly reports of non-fatal suicide attempts based on SPAN data increased between 2012 and 2014, ranging from just over 600 reported attempts in May 2012 to almost 900 in August 2014. VHA’s health care system includes an increasing number of patients with factors, such as a history of suicide attempts, associated with risk for suicide (see Figure 5).

Figure 5: Number of Suicide Attempts Reported Through VA’s Suicide Prevention Applications Network Per Month, 2012–2014




Several steps were taken to assess suicide among VHA users with a history of non-fatal suicide attempt. VHA users with a suicide attempt indication were identified in calendar years 2002–2013 based on indications in inpatient or outpatient encounter records (ICD-9 code E95, excluding E95.9). For each year, the first attempt indication was used as the index date. Non-fatal attempts were identified by survival seven days post-indication date. Among those with non-fatal attempts, suicide and all-cause mortality were assessed from eight to 365 days following the index date of the suicide attempt. As mortality data are currently available only through 2014, analyses are not presented for the 2014 cohort, given limited observable follow-up time. Although both all-cause and suicide-specific 12-month mortality was high among this patient sub-population, all-cause mortality decreased from 2001 to 2014, and suicide rates in the 12 months following attempt remained relatively stable over the period of observation (Figures 6 and 7).

Figure 6. 12-Month All-Cause Mortality Following Suicide Attempt Indication, 2002–2013



**Main Finding: All-cause mortality in the 12 months following a suicide attempt has decreased since 2002.**

Figure 7. 12-Month Suicide Mortality Following Suicide Attempt Indication, by Calendar Year



**Main Finding: Rates of suicide in the 12 months following a suicide attempt have remained stable since 2002.**

#### **D. Total and Gender-Specific VHA Suicide Rates**


For the years 2001–2014, the rate of suicides among all VHA patients who used VHA services in the year of death or in the previous calendar year were evaluated. These are listed in Table 2 and depicted in Figure 8, below. Overall, rates of suicide among all VHA patients decreased between 2001 and 2003, remained relatively stable between 2004 and 2007, and increased between 2008 and 2011. However, a closer examination of rates of suicide among male and female VHA patients indicates that rates of suicide have remained relatively stable among male patients between 2001 and 2014 and increased among female VHA patients during that same time period. Overall, the observed gender-based difference in changes in rates of suicide among female VHA patients between 2001 and 2014 is consistent with comparatively greater increase in rates of suicide among females in the U.S. general population, as reported by CDC earlier this year.

Table 2. Suicide Rates by Sex and Calendar Year

Calendar Year	Suicide Rate (per 100,000 person-years)		
	Total	Males	Females
2001	39.9	42.6	14.4
2002	38.9	41.7	11.7
2003	34.9	37.2	10.7
2004	35.9	38.0	13.3
2005	34.9	36.8	14.7
2006	35.9	38.5	9.0
2007	35.1	37.3	12.5
2008	38.4	40.8	14.3
2009	37.0	39.3	14.8
2010	36.3	38.5	15.4
2011	38.9	41.3	16.3
2012	38.0	40.4	16.0
2013	38.8	41.5	14.4
2014	39.0	41.6	16.7

**Main Finding: Rates of suicide among users of VHA services have remained relatively stable in recent years.**

Figure 8. Suicide Rates Among VHA Users by Sex and Calendar Year



**Main Finding: Rates of suicide among male and female users of VHA services have remained relatively stable in recent years.**

### E. Differences in VHA Patient Suicide Rates by Age and Gender

Table 3 provides information on rates of suicide by age group and gender for VHA patients. In contrast to age-based differences in rates of suicide across age groups in the U.S. general population, although the rates of suicide among those ages 18–29 were lower than or comparable with those of older Veterans in 2001, 18- to 29-year-old VHA patients had the highest suicide rate in 2014 while those 60–79 years of age had the lowest rates that year. It is likely that this finding is strongly influenced by patterns of suicide among males. Among female VHA patients, the highest rates of suicide (2014) were observed for women 40–59 years of age, a pattern that generally held for each year 2001–2014.



Table 3. Suicide Rates by Calendar Year, Overall and by Age and Sex

	CY01	CY02	CY03	CY04	CY05	CY06	CY07	CY08	CY09	CY10	CY11	CY12	CY13	CY14
<b>Total</b>	39.9	38.9	34.9	35.9	34.9	35.9	35.1	38.4	37.0	36.3	38.9	38.0	38.8	39.0
18-29	24.8	30.4	27.0	28.8	21.2	35.5	29.7	36.4	37.7	44.2	50.6	54.2	60.7	58.4
30-39	43.8	41.0	39.5	35.0	36.9	35.8	36.0	33.7	37.2	39.4	44.0	40.5	43.0	45.5
40-49	49.0	47.6	42.5	46.4	44.4	34.2	42.0	42.6	40.1	39.2	45.0	41.2	41.1	40.8
50-59	41.8	42.9	37.9	38.4	36.5	41.1	38.7	43.5	42.1	41.9	45.5	41.1	35.0	39.4
60-69	32.6	29.3	31.1	29.9	29.9	31.7	31.4	36.7	31.5	32.3	30.5	29.2	31.0	32.0
70-79	37.1	35.6	30.3	32.2	31.4	34.2	30.3	32.6	33.5	31.9	32.8	36.3	41.1	33.9
80+	47.5	47.7	36.6	39.9	40.4	36.9	37.8	40.8	41.7	35.4	43.6	44.6	44.0	45.6
<b>Males</b>	42.6	41.7	37.2	38.0	36.8	38.5	37.3	40.8	39.3	38.5	41.3	40.4	41.5	41.6
18-29	33.9	39.5	37.3	38.4	27.8	48.2	37.4	48.0	45.5	55.0	60.4	67.0	74.6	73.3
30-39	54.3	51.2	45.6	42.5	44.7	44.3	43.1	40.0	45.1	47.7	54.3	49.2	51.5	55.0
40-49	55.7	54.7	49.1	52.7	48.6	40.8	49.1	50.6	46.4	44.8	51.8	46.5	46.3	45.6
50-59	44.2	45.8	40.7	40.8	39.3	44.0	41.5	45.9	45.6	45.6	50.2	45.4	39.4	43.4
60-69	33.4	30.5	32.1	30.9	31.1	32.9	32.4	38.0	32.4	32.8	31.1	29.9	32.0	32.9
70-79	38.0	36.2	30.9	32.8	31.5	34.7	30.9	33.1	34.0	32.6	33.3	36.4	42.0	34.3
80+	49.0	49.5	38.1	41.0	41.5	38.0	38.9	41.5	42.9	36.4	44.6	45.7	44.9	46.6
<b>Females</b>	14.4	11.7	10.7	13.3	14.7	9.0	12.5	14.3	14.8	15.4	16.3	16.0	14.4	16.7
18-29	5.7	11.9	5.9	9.0	6.7	4.7	10.2	5.4	15.2	11.1	19.3	11.7	15.7	11.0
30-39	13.1	10.9	21.1	12.6	13.8	10.9	15.7	15.9	14.9	15.3	13.5	14.4	17.4	16.0
40-49	18.1	15.9	13.9	20.5	27.4	7.8	14.8	12.8	16.7	17.4	18.1	20.3	20.6	21.9
50-59	14.5	9.7	5.6	11.2	7.5	12.9	13.8	25.3	18.1	18.8	18.1	18.1	11.8	19.4
60-69	17.1	4.8	9.5	9.6	4.8	6.8	8.6	4.0	9.0	19.4	14.7	11.9	9.7	14.5
70-79	14.6	15.9	7.0	0.0	25.9	5.3	0.0	0.0	10.8	0.0	10.6	31.4	0.0	14.7
80+	14.6	5.9	0.0	16.9	15.3	7.8	8.2	17.7	0.0	0.0	5.2	0.0	5.7	0.0

**Main Finding: Rates of suicide among younger male users of VHA services, ages 18–29, have been rising in more recent years while the suicide rates for other male age groups have remained relatively stable. Rates of suicide among younger female users of VHA services have increased in recent years.**

### **F. Comparison of Suicide Rates among VHA Users and the General U.S. Population**


Table 4 and Figure 9 provide information on changes in relative risk for suicide among VHA patients when compared to members of the U.S. general population. It is important to note that the general U.S. population considers all suicides among U.S. residents, regardless of age or Veteran status. As shown in Table 4, when compared to rates of suicide in the U.S. general population, risk for suicide among all VHA patients, and for males and females separately, has decreased since 2001.

Table 4. Standardized Mortality Ratios Among VHA Health Care Users Compared to the General U.S. Population by Sex and Calendar Year

	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Total	1.65	1.54	1.41	1.46	1.39	1.45	1.39	1.47	1.43	1.38	1.47	1.42	1.42	1.40
Male	1.63	1.53	1.40	1.44	1.37	1.45	1.37	1.46	1.41	1.36	1.45	1.40	1.41	1.38
Female	2.57	2.01	1.83	2.09	2.40	1.41	1.87	2.13	2.14	2.18	2.23	2.12	1.88	2.07

**Main Finding: When compared with the U.S. general population, risk for suicide among users of VHA services has decreased since 2001 for both males and females.**

Figure 9. Standardized Mortality Ratio (SMR) for Suicide Among VHA Users by Sex and Calendar Year



## G. Suicide Among OEF/OIF/OND VHA Users

Risk for suicide following separation from active duty service remains a concern among Veterans of Operation Enduring Freedom, Operation Iraqi Freedom, and Operation New Dawn (OEF/OIF/OND). Rates of suicide among OEF/OIF/OND Veterans who used VHA services are listed by age group and gender in Table 5. Rates of suicide were highest among male OEF/OIF/OND Veterans ages 18–29 years and decreased with age. However, the small number of OEF/OIF/OND Veterans age 60 years and older or among female Veterans limits consideration of age and gender-based differences in risk for suicide among members of this group. Table 6 provides information on rates of suicide among Veterans from active duty service and members of Reserve or National Guard components who were activated in support of operations in Afghanistan and Iraq. Rates of suicide were lower among members of the Reserve and National Guard over the observation period.

Table 5. Suicide Rates Among OEF/OIF/OND VHA Users by Sex, Age Group, and Calendar Year

Sex and Age Group														
	CY01	CY02	CY03	CY04	CY05	CY06	CY07	CY08	CY09	CY10	CY11	CY12	CY13	CY14
<b>Total</b>	0.0	0.0	26.8	26.8	20.6	24.5	30.1	34.3	32.9	30.4	45.2	44.9	50.4	47.4
18-24	0.0	0.0	21.5	54.7	12.5	22.8	35.1	60.5	47.2	46.9	73.0	66.3	85.0	110.3
25-29	0.0	0.0	32.1	37.3	24.6	44.5	33.7	36.5	35.7	38.4	46.9	55.3	67.1	56.3
30-39	0.0	0.0	38.0	7.2	30.6	20.0	35.2	31.7	28.3	28.2	50.0	50.8	52.3	50.5
40-49	0.0	0.0	23.4	8.6	22.3	14.6	21.3	16.5	23.8	17.6	32.5	23.5	31.2	28.0
50-59	0.0	0.0	0.0	0.0	0.0	0.0	7.2	16.7	35.0	10.3	16.3	21.8	12.4	21.4
60-69	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	14.7	0.0	0.0	0.0
70-79	.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
80+	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<b>Male</b>	0.0	0.0	31.6	31.4	21.8	28.2	34.0	38.4	34.7	33.1	49.2	49.1	55.0	52.2
18-24	0.0	0.0	27.0	67.2	15.0	27.0	38.6	69.9	49.7	53.2	75.4	75.3	92.9	124.0
25-29	0.0	0.0	39.5	45.2	29.3	52.4	39.5	41.4	39.5	43.6	52.0	60.9	73.8	62.7
30-39	0.0	0.0	43.9	8.3	35.0	22.8	40.2	36.4	29.9	29.5	57.1	54.6	57.8	56.8
40-49	0.0	0.0	26.5	9.5	14.9	16.3	23.7	16.6	23.6	18.4	33.3	26.4	32.7	28.8
50-59	0.0	0.0	0.0	0.0	0.0	0.0	7.9	18.4	34.0	7.7	18.3	22.1	14.0	22.6
60-69	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	15.5	0.0	0.0	0.0
70-79	.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
80+	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<b>Female</b>	0.0	0.0	0.0	0.0	13.0	0.0	3.4	5.4	20.0	11.1	17.1	16.1	18.8	14.7
18-24	0.0	0.0	0.0	0.0	0.0	0.0	13.8	0.0	29.0	0.0	55.5	0.0	29.1	16.7
25-29	0.0	0.0	0.0	0.0	0.0	0.0	0.0	7.7	12.2	5.2	13.9	17.7	21.5	12.9
30-39	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	17.2	19.8	5.1	27.8	18.9	12.8
40-49	0.0	0.0	0.0	0.0	88.6	0.0	0.0	16.3	25.9	10.6	26.0	0.0	19.0	22.3
50-59	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	44.6	32.9	0.0	19.3	0.0	12.4
60-69	.	.	.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
70-79	.	.	0.0	.	.	.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
80+	.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

**Main Finding: Rates of suicide were highest among younger male OEF/OIF/OND Veterans.**

Table 6. Suicide Rates Per 100,000 Person-Years Among OEF/OIF/OND VHA Users by Active Duty/Reserve Guard by Calendar Year

	CY 07	CY 08	CY 09	CY 10	CY 11	CY 12	CY 13	CY 14
Active	35.0	42.3	36.2	35.4	48.6	48.9	54.9	54.9
Reserve	25.6	25.8	29.1	24.3	40.9	39.4	43.2	34.5

**Main Finding:** When compared with rates of suicide among Veterans of the National Guard or Reserve components, rates of suicide were higher among OEF/OIF/OND active duty Veterans.

## VI. Results – Part 2: Suicide Among All U.S. Veterans, 2001–2014

An important enhancement to this year’s report is the availability of information on rates and characteristics of suicide among all Veterans regardless of VHA use, during the period of observation (2001–2014). Data on suicides among all Veterans were obtained from the VA/DoD Joint Suicide Data Repository. Counts of death for the larger U.S. adult population (18 years of age and older) were obtained from CDC’s WONDER system.<sup>4</sup> Rates of suicide for the civilian population were calculated using estimates of the total U.S. population obtained from WONDER and removing counts for known Veteran suicides for each year within each age and gender subgroup of interest. Crude rates of suicide per 100,000 were calculated for each year and by age and gender for Veterans overall, by use of VHA Services and among civilians. Age adjustment, using the 2000 U.S. standard population, was used to assess differences in rates within groups over time.<sup>5</sup> Estimates of relative risk for suicide were calculated using standardized mortality ratios (SMRs). SMRs can be interpreted as the difference in suicide risk between two populations. SMRs with a value of 1 indicate no difference in risk. SMRs were used to compare rates across groups for any given year, accounting for differences in age and gender composition between the groups.


4. Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death 1999–2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999–2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

5. Klein RJ, Schoenbaum CA. Age adjustment using the 2000 projected U.S. population. Healthy People Statistical Notes, no. 20. Hyattsville, Maryland. National Center for Health Statistics. January 2001.

## A. Magnitude of Veteran Suicide Mortality

In 2001, Veterans accounted for 12.3 percent of the U.S. adult population and 23.1 percent of all suicides among U.S. adults. Between the years 2001–2014, both the proportion of U.S. adults who were Veterans (8.5 percent, 2014) and the proportion of adult suicide decedents who were Veterans decreased (17.9 percent, 2014). However, changes in the proportion of U.S. adults who were Veterans or the proportion of adult suicides who were Veterans leaves gaps in our understanding of changes in rates of suicide among Veterans over time. Therefore, steps were taken to test for changes in rates of suicide among Veterans and control for shifts in the demographic composition of populations over time. Age-adjusted rates of suicide were calculated for each year during the study period using the 2000 U.S. standard population weights. As shown in Figure 10, differences in age-adjusted rates of suicide were greater for Veterans than civilians between 2001 and 2014, with substantial increases observed for female Veterans between years. Further differences within Veteran subpopulations were observed when changes in the age-adjusted rates of suicide were calculated separately for Veterans with and without use of VHA services. Overall, Non-VHA Veterans had greater increases in rates of suicide when compared to changes in rates of suicide among VHA Veterans. Further, females Veterans were the only group with a decrease in rate of suicide over time.


Figure 10. Percent Changes in Age-Adjusted Rates of Suicide Among Veterans and Civilians, 2001–2014



## Average Number of Veteran and Civilian Suicides per Day


We calculated the average number of Veterans and Civilian Suicides per day by taking the total number of suicides in each group for each year and dividing by 365, the number of days in a year. In 2001, an average of 19 Veterans died per day by suicide (Figure 11). This number increased slightly from 2001 to a high of 21 per day in 2010, with a subsequent decrease to 20 per day in 2011 and remaining stable since that time. In contrast, the average number of civilian adults who died by suicide each day has increased steadily from 62 per day in 2001 to 93 per day in 2014. When considering VHA service use prior to death, the average number of VHA Veterans who died by suicide each day has increased from 4 in 2001 to 6 in 2014 (Figure 12). It should be noted that decreases in the size of the Veteran population and contrasting increases in the size of the U.S. population limit interpretation of these statistics. Rates of suicide, stratified by group, are more appropriate for understanding changes in risk among Veterans and civilians and are provided throughout this report.

Figure 11. Average Number of Suicides Per Day Among Veterans and Civilians by Year, 2001–2014



**Main Finding: On average, the number of civilians who died by suicide each day has increased each year since 2001.**

Figure 12. Average Number of Suicides Per Day Among Veterans With and Without Use of VHA Services, 2001–2014




**Main Finding: On average, the number of Veterans who died by suicide each day has remained stable since 2011.**

**B. Comparison of Suicide Rates Among Veterans Who Do and Do Not Use VHA Services, 2001–2014**

In 2014, male VHA Veterans were only 24 percent more likely to die by suicide than male non-VHA Veterans. Similarly, excess suicide risk among VHA female Veterans decreased from 86 percent to 5 percent between 2001 and 2014. Since 2013, there has been no statistically significant difference in risk for suicide among female Veterans with and without use of VHA services.

Figure 13. Standardized Mortality Ratios (SMR) Comparing Risk for Suicide Among Veterans With and Without Use of VHA Services by Calendar Year and Sex, 2001–2014




**Main Finding: The difference in suicide risk between Veterans with and without the use of VHA services has diminished since 2001.**

### C. Comparison of Veteran and Adult Civilian Suicide Risk, 2001–2014

Prior to 2006, Veteran suicide rates were lower than adult civilian suicide rates after accounting for age and gender differences between the populations. Risk of suicide among Veterans relative to civilians has increased relatively steadily since 2001. In 2014, Veterans were 21 percent more likely to die by suicide when compared to their adult civilian peers, adjusting for age and gender. Differences in estimates of relative risk were observed for Veterans with and without use of VHA services prior to death, with VHA Veterans experiencing higher suicide rates than adult civilians across the observed time period (Figure 14). A greater increase in the relative risk for suicide among Veterans when compared to civilians was observed for those without use of VHA services over this time period.



Figure 14. Standardized Mortality Ratios for Veterans With and Without Use of VHA Services Compared to Civilians, 2001–2014




**Main Finding: When compared with suicide mortality among the civilian population, a greater increase in the relative risk for suicide among Veterans was observed for those without use of VHA services.**

### Age Differences in the Comparison of Veteran and Civilian Suicide Risk


Figures 15 through 21 provide age-specific suicide rates for all Veterans and civilians, by year. Figures 22 through 24 present this information for males, and Figures 25 through 27 present this information for females. Overall, rates of suicide have increased more for Veterans than among their civilian peers. However, there are important differences across age groups and gender. For example, rates of suicide have remained relatively stable for Veterans ages 40–49, and little difference was observed in rates of suicide among older female Veterans when compared to similarly aged civilian females. However, the comparatively small number of suicides among older female Veterans limits consideration of observed differences.

Figure 15. Crude Rates of Suicide by Calendar Year Among Veterans (V) and Civilians (C) Ages 18–29 Years, 2001–2014




**Main Finding: Rates of suicide have increased substantially among younger Veterans while remaining relatively stable among civilians 18–29 years of age.**

Figure 16. Crude Rates of Suicide by Calendar Year Among Veterans (V) and Civilians (C) Ages 30–39 Years, 2001–2014




**Main Finding: Rates of suicide have increased among Veterans ages 30–39 while remaining relatively stable among civilians in this age group.**

Figure 17. Crude Rates of Suicide by Calendar Year Among Veterans (V) and Civilians (C) Ages 40–49 Years, 2001–2014




**Main Finding: Despite an increase in suicide rates among the civilian population ages 40–49, rates of suicide have remained stable among Veterans in the same age group.**

Figure 18. Crude Rates of Suicide by Calendar Year Among Veterans (V) and Civilians (C) Ages 50–59 Years, 2001–2014




**Main Finding: Rates of suicide increased substantially among Veterans ages 50–59. Increases in civilian suicide rates are also evident in this age group.**

Figure 19. Crude Rates of Suicide by Calendar Year Among Veterans (V) and Civilians (C) Ages 60–69 Years, 2001–2014




**Main Finding: Rates of suicide increased substantially among Veterans ages 60–69. Increases in civilian suicide rates are also evident in this age group.**

Figure 20. Crude Rates of Suicide by Calendar Year among Veterans (V) and Civilians (C) Age 70–79 Years, 2001–2014




**Main Finding: Despite increases in suicide rates among the civilian population ages 70–79, rates of suicide remained stable among Veterans in the same age group.**

Figure 21. Crude Rates of Suicide by Calendar Year Among Veterans (V) and Civilians (C) Age 80+ Years, 2001–2014



**Main Finding: Rates of suicide increased among Veterans age 80 and older while remaining stable among civilians in this age group.**


Figure 22. Crude Rates of Suicide by Calendar Year Among Male Veterans (V) and Civilians (C) Ages 18–39 Years, 2001–2014



**Main Finding: When compared with civilian males, suicide rates were higher among younger male Veterans with greater and increases to a greater degree over time.**




Figure 23. Crude Rates of Suicide by Calendar Year Among Male Veterans (V) and Civilians (C) Ages 40–69 Years, 2001–2014




**Main Finding: Increases in rates of suicide among male Veterans ages 50–69 were larger than those observed among civilian males in the same age groups.**

Figure 24. Crude Rates of Suicide by Calendar Year Among Male Veterans (V) and Civilians (C) Age 70+ Years, 2001–2014




**Main Finding: Rates of suicide among older adult male Veterans were lower than rates of suicide among older adult civilian males across the time period.**

Figure 25. Crude Rates of Suicide by Calendar Year Among Female Veterans (V) and Civilians (C) Ages 18–39 Years, 2001–2014




**Main Finding: Greater increases in rates of suicide increased to a greater degree were observed among younger female Veterans than among younger female civilians.**

Figure 26. Crude Rates of Suicide by Calendar Year Among Female Veterans (V) and Civilians (C) Ages 40–69 Years, 2001–2014



**Main Finding:** Rates of suicide were higher among female Veterans ages 40–69 when compared with suicide rates among female civilians in the same age groups.

Figure 27. Crude Rates of Suicide by Calendar Year Among Female Veterans (V) and Civilians (C) Age 70+ Years, 2001–2014




**Main Finding: Despite relative instability associated with a comparatively smaller number of older female Veterans, rates of suicide among older female Veterans were similar to rates of suicide among older adult female civilians.**

## D. Mechanisms of Veteran and Civilian Suicide, 2001–2014

Use of firearms is associated with the highest rate of suicide mortality compared with other prevalent suicide mechanisms in the U.S. Figures 28 and 29 demonstrate that the proportion of suicide decedents using firearms is higher among both male and female Veterans than among the adult civilian population. In addition, while the proportion of civilian decedents who used firearms has decreased, it has remained relatively stable or increased slightly for both male and female Veterans.


Figure 28. Civilian Suicide Deaths by Mechanism and Sex, 2001–2014



**Main Finding: From 2001 to 2014, the percentage of suicides resulting from a firearm injury decreased among U.S. adult civilians, for both males and females.**


Figures 30 and 31 show the proportion of all deaths by mechanism among all Veterans and among Veterans with and without use of VHA services in 2001 and 2014. In contrast to trends in the U.S. civilian population, the proportion of suicides resulting from a firearm injury has increased among female Veterans and has remained relatively constant among males. Among female Veterans, there were decreases in the proportion of suicides resulting from poisoning and increases in the proportion of suicides resulting from suffocation and firearms. The observed increase in suicides resulting from suffocation was greater among female Veterans who used VHA services than among female Veterans without VHA service use (Figures 30 and 31).

**Figure 29. All Veteran Suicide Deaths by Mechanism and Gender in 2001 and 2014**



**Main Finding: The percentage of all suicides resulting from a firearm injury remained high among Veterans from 2001 to 2014.**


Figure 30. VHA Veteran Suicide Deaths by Mechanism and Gender, 2001–2014



**Main Finding: The percentage of all suicides resulting from suffocation and firearms increased among female Veterans who used VHA services.**



Figure 31. Non-VHA Veteran Suicide Deaths by Mechanism and Gender in 2001 and 2014



**Main Finding: The percentage of suicides resulting from a firearm injury was similar among Veterans with and without the use of VHA services.**

### E. Understanding the Burden of Veteran Suicide: Magnitude vs. Risk

When directing suicide prevention efforts it's important to consider the distribution of suicides as well as differences in rates among key population subgroups. Figures 32 through 35 demonstrate important differences in the distribution of the number and rate of suicide across age groups and gender when compared to the characteristics of suicide among civilians. As demonstrated in Figures 32 and 33, rates of suicide are highest among younger male Veterans and lowest among male Veterans ages 60–79 years. However, the greatest number of suicides among male Veterans was observed for those ages 50–69 years. In contrast, the greatest number of suicides among civilian males was observed for those ages 59 and younger. Similarly, while fewer differences in the distribution of counts of suicide between civilians and Veterans are observed for women, the greatest rates of suicide among female Veterans were among those between 18 and 29 years of age.

Figure 32. Comparison of Suicide Counts and Rates by Age Group for Civilian Males, 2014





Figure 33. Comparison of Suicide Counts and Rates by Age Group for Veteran Males, 2014



**Main Finding:** Among civilian males, the largest number of lives lost to suicide occurs among younger and middle-aged adults (ages 18–59) with the highest rates of suicide among older adults. Among male Veterans, the largest number of lives lost to suicide are middle-aged (ages 50–69), with the highest rates of suicide are among the youngest males (ages 18–29).

Figure 34. Comparison of Suicide Counts and Rates by Age Group for Civilian Females, 2014





Figure 35. Comparison of Suicide Counts and Rates by Age Group for Veteran Females, 2014



**Main Finding:** Among civilian females, age-specific suicide rates correspond closely with the number of lives lost to suicide, with both peaking between the ages of 40–59 years. Among Veteran females, the largest number of lives lost to suicide occurs in middle age (ages 40–59), but the highest rate occurs among female Veterans ages 18–29.

## VII. Summary and Discussion of Findings

This comprehensive analysis of Veteran suicide from 2001 to 2014 confirms that when compared to their non-Veteran peers, most Veterans are at an increased risk for suicide. However, important differences in rates of suicide among Veteran and civilian groups were observed when rates were stratified by sex and age group. While the average number of Veterans who died by suicide each day has remained relatively stable in recent years, the relative risk for suicide among Veterans when compared to civilian adults has increased, as have age-specific suicide rates for Veterans 18–39, 50–69, and 80+ years of age. Increases in suicide rates are particularly evident among female Veterans and Veterans who do not use VHA services.

Despite evidence of increases in suicide rates among most civilian and Veteran groups, rates of suicide have remained stable among some Veteran sub-populations, such as males ages 40–49, and differences have diminished between Veterans with use of VHA services.

The most common means for suicide among Veterans is firearms, with approximately 41 percent of female and 68 percent of male Veteran suicide deaths resulting from a firearm injury in 2014. Poison is the second-most common means of suicide for female Veterans—32 percent of female Veterans who die by suicide use poison. Among male suicide decedents, suffocation is the second-most common cause of death (17 percent, 2014). The use of firearms as the mechanism for suicide death decreased among civilians from 2001 to 2014, but remained stable among Veterans. These results strongly suggest that firearms safety initiatives are likely an important component of an effective suicide prevention strategy for male and female Veterans.

Among male Veterans, suicide rates are highest in the younger and older years, and among female Veterans, suicide rates are highest in the younger years. However, because the age distribution of the living Veteran population is heavily weighted toward middle-aged adults, the resulting burden of suicide, in terms of the number of lives lost, is most evidenced among middle-aged Veterans, despite the lower rates of suicide observed for this sub-population. Analysis of suicide risk within the Veteran population indicates that it is important to develop and direct Veteran suicide prevention initiatives that are tailored to reach Veterans of all ages.

Findings included in this report highlight the complex relationships between a history of U.S. military service and suicide risk and the need for additional assessments to understand factors such as changes in risk exposures, workforce composition and socioeconomic factors, and rates of suicide across Veteran and civilian groups.

## VIII. Suicide Data Report Part II

The findings presented in this report represent the first analysis of more than 50 million Veteran suicide records. This epidemiologic overview of suicide risk among the Veteran population and how it compared to suicide risk over time among U.S. adult civilians sets the stage for continued analyses. Additional analyses of this data are ongoing and will include the following inquiries, at a minimum.

- Comparison of suicide risk and burden among VHA and non-VHA OEF/OIF/OND Veterans
- Comparison of rates of suicide among residents in urban and rural areas
- Rates of suicide across branches of military service (i.e., Army, Navy, Air Force, Marines)
- Rates of suicide by race/ethnicity
- Risk for suicide during periods of transition

### *References:*

\* Source: VA Benefits & Health Care Utilization Pocket Card, Updated 5/13/16; Veteran Population as of 09/30/15 (<http://www.va.gov/vetdata/docs/pocketcards/fy2016q3.pdf>)

\*\*Source: VA Benefits & Health Care Utilization Pocket Card, Updated 5/13/16; Produced by the National Center for Veterans Analysis and Statistics. (<http://www.va.gov/vetdata/docs/pocketcards/fy2016q3.pdf>). (pg 1)

Curtin SC, Warner M, Geedegaard H. Increase in suicide in the United States, 1999–2014. NCHS data brief, no, 241. Hyattsville, MD: National Center for Health Statistics. 2016. Details on priority group enrollment criteria can be found at:

[http://www.va.gov/HEALTHBENEFITS/resources/priority\\_groups.asp](http://www.va.gov/HEALTHBENEFITS/resources/priority_groups.asp) (pg 6)

Harris, E.C. & Barraclough, B. (1997) Suicide as an outcome for mental health disorders. A meta-analysis. Br J Psychiatry, 170, 205-228 (pg 8)